

Coda Q tomography at the Galeras volcano, Colombia

Carlos A. Vargas Jiménez⁽¹⁾, *Juan P. Duran*⁽¹⁾
and Lluís G. Pujades⁽²⁾

(1) *Department of Geosciences, Universidad Nacional de Colombia, Bogotá, Colombia*

(2) *Technical University of Catalunya, Barcelona, Spain*

Abstract

Spatial variation of coda Q (Q_c) was analyzed in the Galeras volcano, located in the Andes of southwestern Colombia. The Q_c value was calculated using the Single Backscattering (SBS) model working with seismograms filtered in the band centered on 3.0 Hz. We used the estimated values as the mean attenuation that represents the total effect of simple scattering in an ellipsoidal volume. Then an inversion procedure was implemented to estimate the spatial variation of Q_c as function of volume fractions. Synthetic tests in 2D and 3D allow us to extend the procedure with available data from the seismological network located at the Galeras volcano.

The inversion of 2855 Q_c values in a grid of 8x8x8 nodes associated to 435 earthquakes and 19 stations permitted to resolve the presence of two levels of attenuation anomalies which are assumed to be related to bodies of partially melted rocks located respectively at 9.0 km and more than 13.3 km from the summit. These bodies could be responsible for the vulcanian type explosions as a consequence of mixture of shallow magma with deep basic magma. This shallow body seems to extend to the summit and may represent the presence of plugs and fluids that are constantly fed from the deeper body and produce the actual hydrothermal activity with eruptive events. A recent event at November, 2005, generated the evacuation of people (aprox. 9000 persons) of the at least four towns around the Galeras volcano (Mapachico, Genoy, La Florida and Nariño). However, the big volume involved in this process suggests that more eruptive events may be expected at the Galeras volcano.

1. Introduction

The Galeras volcano (lat. 1°14' N, lon. 77°22' W) is a 4200m-high andesitic volcano in the Andes of southwestern Colombia (Nariño department) and it is the most recent active center of the Galeras volcanic complex (Cepeda, 1986). The cone of this active volcano has a 320m diameter and it lies within a caldera that is breached to the west. The Galeras volcanic complex is historically characterized by vulcanian type explosions (Calvache, 1990), the most important one, recorded in 1936, generated a pyroclastic flow to the northeast of the volcanic structure. The volcano lies 8 km from the Pasto city (Figure 1) and various other towns are located on its flanks. In total, about 400.000 people live close to the volcano, inside the hazardous zone.

Since 1988, when the Galeras volcano was reactivated, after 50 years of calm, it entered in a fumarolic and degassing stage, with seven eruptions between 1989 and 1993. An andesitic dome was extruded in September 1991 and was destroyed during the July 1992 eruption. Since 1994 the Galeras is in a quiet stage with lesser explosions, and seismic crises in the years of 1995, 1998, 2000, 2002, 2004, 2005, 2006, 2008 and 2009.

Few studies have been performed in the Galeras volcano. The first works were oriented to describe the evolution and its geological frame (Cepeda, 1986; Calvache, 1990). Because in beginnings of the year's 1990 Galeras volcano was declared the Decade Volcano, several works were done in order to propose its eruptive mechanism throughout geological, geochemical and geophysical observations. Some of these studies were based on the assumption of two magmatic deposits (Calvache and Williams, 1992; Muñoz et al., 1993; Stix et al., 1993; Fisher et al., 1996; Calvache and Williams, 1997a,b; Calvache et al., 1997; Banks et al., 1997; Fisher et al., 1997; Stix et al., 1997; Sano et al., 1997; Zapata et al., 1997). In the seismological field, Gomez and Torres (1997) gave the first study related with the coda wave attenuation in low-frequency volcanic seismic events. On the other hand, Narvaéz et al. (1997), Gomez et al. (1999) and Seidl et al. (1999) have worked with the screw-type signals and the features of the different seismic sources. However, nobody has worked on the spatial variation of the attenuation, so this is an innovative study that enlarges the knowledge about the Galeras volcano.




Figure 1. Location of the Galeras volcanic complex in the Nariño department, Colombia. Triangles represent the seismological stations used in this study.

2. Geotectonic setting

The Galeras volcano is in the Galeras Volcanic Complex (GVC), located in southwestern Colombia, a region with a metamorphic rocks basement of Precambrian and Paleozoic ages. The basement is overlaid by metamorphic rocks of Cretaceous age with low and medium grade associated with amygdular metabasalts. All this is covered by volcano-sedimentary units of Tertiary age that made a plateau, over which the Pleistocenic and Holocenic volcanoes have emerged (Hurtado and Cortés, 1997). The GVC has erupted lavas and pyroclastic flows, ranging from basaltic andesites to dacites, during the last million years. The chemical composition of the volcanic products has been fairly constant, characterized by calc-alkaline, high-silica, medium-potassium andesites. Variations from basaltic andesites to dacites have occurred during individual stages, but overall, composition and petrographic characteristics are quite constant (Calvache and Williams, 1997a).

The tectonic of this region is very complex, being the result of the collision between the Nazca and South American plates, which causes the uplift of the Andes and the volcanism in the region. The structural trend is N40°E and the principal tectonic feature is the Romeral Fault Zone, which has been interpreted as the limit between continental crust at East and the oceanic crust at West (Barrero, 1979). This system includes the Silvia-Pijao and Buesaco faults. Both faults cut the Galeras volcano structure, and are associated with many old caldera systems. The Machabajoy, Pasto, Guitara, San Ignacio and Magdalena faults limit the volcano structure with similar trend (Figure 1).

3. The data

The seismic data used in this study covered 435 earthquakes recorded in the period between September 1989 and June 2002. The seismograph network installed on the Galeras is composed of 19 stations equipped with vertical component L-4C seismometers with 1.0 second natural period. All seismograph stations, however, have not operated continuously through time because various modifications in the




Figure 2. Typical seismogram of a local earthquake occurred in the Galeras volcano and analyzed in this work. The event was a $ML=1.8$, and was located at Lat N $1^{\circ} 15.27'$, Lon W $77^{\circ} 19.74'$ and Depth= 8.18 km.

network configuration. The ground motions are sent by radio to an observatory in Pasto city and the signals are sampled with an ADC system of 12 bits at a rate of 100 s⁻¹. The complete system has a flat response between 1.0 and 25.0 Hz. We used earthquakes with local magnitude less than 2.0 to avoid clipping. Figure 2 shows a typical seismogram of a local earthquake occurred in the Galeras volcano and analyzed in this work.

3. Method of Analysis

3.1 Coda Q

We follow the Single Backscattering (SBS) model of Aki and Chouet (1975) for calculating the coda Q^{-1} . This assumes that the coda of a local earthquake is composed of the sum of secondary S waves produced by heterogeneities distributed randomly and uniformly in the lithosphere. The coda is the portion of seismogram corresponding to back-scattered S-waves. We have estimated Q_c^{-1} values using the following equation (Aki and Chouet, 1975):

$$P(\omega, t) = \frac{2g(\theta)|S(\omega)|^2}{\beta t^2} e^{\left(\frac{-\omega t}{Q_c}\right)} \quad (1)$$

where, $P(\omega, t)$ is the time-dependent coda power spectrum, ω is the angular frequency, β is the shear wave velocity, $|S(\omega)|$ is the source spectrum, and $g(\theta)$ represents the directional scattering coefficient, defined as 4π times the fractional loss of energy by scattering per unit travel distance of primary waves and per unit solid angle at the radiation direction θ measured from the direction of primary wave propagation. Under these considerations, the geometrical spreading is assumed to be proportional to r^{-1} , which only applies to body waves in a uniform medium. On the other hand, the source factor can be treated as a constant value for single frequency, according to equation (1) Q_c^{-1} values can be obtained as the slope of the least squares fit of $\ln [t^2 P(\omega, t)]$ versus ωt , for $t > t_\beta$, where t_β represents the S-wave travel time (Rautian and Khalturin, 1978).

3.2 Regionalization technique: a 3D approach

Following Singh and Herrmann (1983), a way to regionalize for Q_c is based on the work of Malin (1978) who, expanding on the work of Aki (1969) and Aki and Chouet (1975), realized that the first order scatterers responsible for the generation of coda waves at a given lapse time are located on the surface of an ellipsoid having earthquake and station locations as foci. According to Pulli (1984), in the ellip-

soidal volume sampled by coda waves at a time t , the large semi-axis is defined as $a_1 = \beta t/2$, and the small semi-axis as $a_2 = a_3 = (a_1^2 - r^2/4)^{1/2}$, where r is the source-receiver distance of the ellipsoid. We have confirmed the volume sampled by coda waves in the case of single isotropic scattering, simulating numerically the movement of many particles, which represents the propagation of S-wave energy using a Monte Carlo method.

The basic assumptions for the numerical simulation are:

- (1) the scattering is isotropic;
- (2) scattered waves are incoherent and point-like scatterers are randomly distributed with a uniform density;
- (3) a short duration point-like source radiates S waves spherically;
- (4) there are no conversions between P and S waves, which travel with a constant S wave velocity.

The assumptions and procedure are described in Hoshiya (1991). Figure 3 shows the simulation test for the case of single scattering. It can be observed that, in the first order scattering the sampled volume in the attenuation process is closer to the segment defined by the hypocenter-station path that delineates an ellipsoid with focus hypocenter (F_1) and station (F_2). Consequently, we have assumed that the average sampled volume can be represented by $t_{av} = t_{start} + \Delta t_{win} / 2$, where t_{av} is the average lapse time for the first order scattering, t_{start} is the starting lapse




Figure 3. Monte Carlo simulation of the volume sampled in the single scattering process. For the simulation, an arbitrary coordinate system centered at $(0, 0, 0)$ km was chosen, with an hypocenter (F_1) located at $(-200, -200, 0)$ km and one station (F_2) at $(200, 200, 0)$ km. A lapse time of 100 s and a velocity of 3.0 km/s were considered. (a) Horizontal projection of the Monte Carlo simulation. (b) 3D perspective of the Monte Carlo simulation.

time, and Δt_{win} is the window length. Then, by knowing the volumes of the ellipsoids as a function of time, a generalized inversion for regional Q_c can be performed. For the purpose of inversion, we define a geographic grid about the seismic station that also encloses the hypocenter. Herein, we recognize that each measured Q_c is an average estimate Q_{av} (or Q_{apparent}) for the volume sampled by the ellipsoid given by:

$$\frac{V_{\text{TOTAL}}}{Q_{\text{av}}} = \sum_j \frac{V_{\text{Block}.j}}{Q_j} \quad (2)$$

where $V_{\text{Block}.j}$ is the fraction of volume (block) in an ellipsoidal region with the true attenuation coefficient Q_j (or Q_{true}). Assuming a constant velocity of propagation, the volume travelled by a ray that leaves the hypocenter, moves out to the ellipsoid defined by the observation time of the coda, and is scattered to the receiver, can be determined.

Equation (2) can be written as:

$$\frac{1}{Q_{\text{av}}} = \frac{1}{Q_1} \frac{V_{\text{Block}-1}}{V_{\text{TOTAL}}} + \dots + \frac{1}{Q_1} \frac{V_{\text{Block}-j}}{V_{\text{TOTAL}}} + \dots + \frac{1}{Q_n} \frac{V_{\text{Block}-n}}{V_{\text{TOTAL}}} \quad (3)$$

where, the ratio $V_{\text{Block}.j}/V_{\text{TOTAL}}$ is just the volume fraction associated to the total scattered wave travel path spent in the j^{th} block. If the process is repeated for each station-hypocenter pair, the entire region is sampled. Equation (3) is of the form

$$a_1 x_1 + \dots + a_i x_i + \dots + a_n x_n = y \quad (4)$$

where $y = \left(\frac{1}{Q_{\text{av}}} \right)$, $x_i = \left(\frac{1}{Q_i} \right)$, $a_i = \left(\frac{V_{\text{Block}-i}}{V_{\text{TOTAL}}} \right)$.

Then, a least squares estimate of the x_i is given by the compact matrix equation:

$$\mathbf{AX} = \mathbf{Y} \quad (5)$$

where \mathbf{A} is a $(k \times n)$ coefficient matrix, \mathbf{X} a $(n \times 1)$ vector, \mathbf{Y} a $(k \times 1)$ vector and k is the number of station-hypocenter pairs. A solution of (5) is estimated using the damped least square technique of Crosson (1976). According to this technique the solution and resolution matrices can be found with next equations:

$$\mathbf{X} = \left[\mathbf{A}^T \mathbf{A} + \sigma^2 \mathbf{I} \right]^{-1} \mathbf{A}^T \mathbf{Y} \quad (6)$$

$$\mathbf{R} = \left[\mathbf{A}^T \mathbf{A} + \sigma^2 \mathbf{I} \right]^{-1} \mathbf{A}^T \mathbf{A} \quad (7)$$

The method described above was used for the regionalization of Q_c in several synthetic tests in the Galeras volcano.

3.3 Synthetic tests

Several tests were done to evaluate the efficiency of this procedure. First we solved the problem in 2D for a distribution of regular attenuation anomalies in an area where many station-epicenter pairs were well dispersed. Figure 4.a shows a model with two areas associating different attenuation values that are divided into 8×8 cells. In this case only two Q_c values have been used namely $Q_c = 30$ and $Q_c = 90$. On this attenuation distribution 500 random ellipses were calculated, that allowed sampling the model mainly in the central zone (Figure 4.b). Then we calculated elliptical areas and its Q_{av} (pondered by the distribution attenuation areas) associated with every hypothetical station-epicenter pair. The area fractions in a grid of regular cells permitted building an equation system as:




Figure 4. 2D synthetic test in a grid of 8×8 cells for evaluating the inversion procedure. (a) Original attenuation distribution associated to two different areas. (b) Random ellipses with variable foci distances (earthquake – station) that have sampled the area mainly in the central zone. (c) Inversion of attenuation. (d) Frequency of sampled cells (EPC). (e) Distribution of RDE.

$$\frac{A_{\text{TOTAL}}}{Q_{\text{av}}} = \sum_j \frac{A_{\text{Cell},j}}{Q_j} \quad (8)$$

Figures 4.c, 4.d and 4.e represent respectively the inverted model, the number of Ellipses per Cell (EPC) and the resolution diagonal elements matrix (RDE) of equation (8). As we can see, in cells covered with enough ellipses the solution is reliable and true attenuation values ($1/Q_c$) were well restored. A second 2D synthetic test was developed to evaluate the performance in a more complex frame using a chessboard (Figure 5.a). Again, we can see that contrast between high and low attenuation values were well restored (figure 5.b to 5.e). However, these results suggest that there are necessary many station-epicenter pairs with different r that guarantees a well covered area.




Figure 5. Synthetic test in a chessboard of 8x8 cells. (a) Original attenuation distribution associated to two different values. (b) Random ellipses with variable foci distances. (c) Inversion of attenuation. (d) Frequency of sampled cells (EPC). (e) Distribution of RDE.

Similarly, we have extended this procedure to 3D models. Two synthetic tests have been analyzed. The first one we have considered is a 3D chessboard with Q_c values of 30 and 90. In this scenario we have considered 2000 ellipsoidal foci that sampled well the entire volume (figure 6.a). We have supposed an ideal frame where stations can be located in depth, in places that suggest us to have a better volume sampled. Figure 6.b shows how the inversion procedure, in a grid of $8 \times 8 \times 8$ blocks, highlights reasonably the attenuation values in the volumes well sampled. Similarly, the RDE and Ellipsoids per Cell (EPC) values represents a good tool to detect the zones well solved and covered with enough ellipsoids (Figure 6.c).

However, we wished to know how is the behavior of available data (earthquakes and stations) to determine the presence of attenuation anomalies. A new test was considered to evaluate the inversion performance to determine the same structure used in last test with 19 stations and 435 earthquakes. Figure 7.a shows distribution of stations and earthquakes located mainly between 0.0 and 10.0 km depth from summit of the Galeras volcano. The dark points in the below of Figure 7.a represents nodes associated to a grid of $8 \times 8 \times 8$ cells that has been used to inversion of Q_c values. This distribution of foci allowed us to have sampled volumes that cover almost all the Galeras volcano structure. As we can see in Figure 7.b, a comparison of 3D Q_c values distribution between original and inverted model suggests a well solved model mainly where there are enough concentration of ellipsoids, that is, in the longitude range between 77.5°W and 77.2°W ; latitude range between 1.1°N and 1.5°N , and depth range located between 5.5 and -16.7 km. However, the distribution of RDE and EPC values shows that above 5.5 km and below -11.0 km the results would not be reliable.




Figure 6 (a)




Figure 6 (b)




Figure 6 (c)

Figure 6. Synthetic test in a 3D attenuation model of $8 \times 8 \times 8$ blocks.

- (a) Distribution of 2000 random pairs of foci (left) and ellipsoids related to volume sampled in the simple scattering process (right).
- (b) Original and inverted attenuation models with Q_c values of 30 and 90.
- (c) Spatial distribution of RDE and EPC.




Figure 7 (a)




Figure 7 (b)




Figure 7 (c)

Figure 7. Synthetic test where a 3D attenuation model has been simulated with real information of foci pairs (earthquake-station).

- (a) Distribution of 435 earthquakes and 19 stations in the Galeras volcano zone (up), and profiles N-S and E-W of the ellipsoids related to volume sampled in the simple scattering process (below). The dark points represent distribution of nodes associated to grid of 8x8x8 blocks. The ellipsoids have been cut by the intersection with the surface
- (b) Original and inverted attenuation models with Q_c values of 30 and 90.
- (c) Spatial distribution of RDE and EPC.

4. Data analysis and results

The 435 earthquakes selected for this study permitted to calculate 2855 Q_c values. Initially, for hypocenter determinations we used the computer program HYPO71 (Lee and Lahr, 1985). We used the velocity model adopted by the Pasto Volcano Observatory (Table 1), defined using regional seismic reflection profiles and based in the work by Meissner et. al. (1976). We carefully selected seismograms of located earthquakes with root mean square residuals of calculated travel times and observed times (RMS) less than 0.1s, azimuthal separation between stations (GAP) less than 180° , and formally computed horizontal and vertical errors in locations of less than 1 km. Figure 7.a shows the location of the seismograph stations and selected earthquakes used where we can see a range depth between 0.0 and 10 km.

Table 1. *Velocity model used for locate the hypocenters. This model is used by the Seismological and Volcanological Observatory at Pasto city (INGEOMINAS), which was derived from the seismic profiles developed for Meissner et. al. (1976).*


Depth at top of the layer (km)	P-wave velocity (km/s)
0.0	3.50
2.0	3.70
4.0	4.00
8.0	6.00
26.0	6.80
44.0	8.00

4.1 Coda Q


The time-dependent coda power spectrum, over a time-window of 2.0s, was calculated as the mean squared amplitudes of the coda $A_{\text{obs}}(f|r, t)$ from band pass filtered seismograms around centre frequency 3.0 Hz with the respective bandwidth 1.0–5.0 Hz. The squared amplitudes were averaged in a 50% overlapping time window of $t \pm 0.2$ s. Then, Q_c^{-1} was estimated for each seismogram by means of a least-squares regression of equation (1) using a time window of $2(r/\beta)$ to $2(r/\beta)+5$ s from the origin time of the earthquake. Following Aki and Chouet (1975) we have estimated a pondered average S wave velocity of $\beta= 3.0$ km/s in the depth range 0 to 26 km, assuming a $V_p/V_s = 1.78$ (Moncayo et al., 2004), which permit to find the Q_c^{-1} values and to extend them to the tomographic procedure under the hypothesis SBS model and the assumptions of Hoshiaba (1991). Only those amplitudes higher than twice the background noise level were considered for the regression and only Q_c^{-1} values obtained from least-squares fit with correlation coefficients greater than 0.7 were kept. With the central frequency and the β selected we can estimate anomalies with a minimum side of 1 km.

4.2 3D coda Q structure under the Galeras volcan

The dense data coverage at Galeras volcano allow to find a zone resolved in the depth range between 3.3 km to -13.3 km, longitude range -77.32° to -77.51° , and latitude range 1.15° to 1.36° (Figure 8.a). This volume is associated to several attenuation anomalies that suggest the presence of an intricate system of magma bodies. At least three main attenuation anomalies are evident, at southeast, at southwest and at northwest of the main cone (Figure 8.b). These possible bodies have



(a)



(b)

Figure 8. Distribution of Q_c values in Galeras volcano inverted from 435 earthquakes and 19 stations. (a) Spatial distribution of RDE and EPC. (b) Attenuation distribution in depth by mean eight sections through the Galeras volcano.

geometric centres at depths of 9.0 km at NW, >13.3 km at SW, and > 13.3 km at SE from crater and have connections through several minor anomalies that, although they are not totally clear in depth, could be associated with the Silvia-Pijao and Buesaco faults. In general, we can see that the rock surrounding the anomalies has Q_c values between 50 and 100, and the anomalies have contrast Q_c values between 10 and 30.

5. Discussion

Although in this study we did not have discriminated intrinsic and scattering attenuation, Vargas *et al.* (2004) reported that 50 km at northeast of this zone (CRU station) the contribution in the coda wave attenuation at $f=3$ Hz is equally associated to scattering and intrinsic effects. It means that the seismic albedo close to 0.5 represents heterogeneity contributions for the scale length of the analyzed frequency that could be related to the tectonic setting that allow develop the volcanoes in the GVC. Sanchez *et al.* (2005) reported two- and three-dimensional mapping of b-value that illuminates a vertically elongated structure beneath the active crater of Galeras, which may be associated with a conduit, or alternatively, with a shallow region of temporary magma storage. Any case could be related to the fracture system that allows the magma emplacement and explains the scattering effects identified in the attenuation analysis.

On the other hand, it seems very feasible that the intrinsic attenuation contributions could be associated to rocks partially melted or rich in fluids. Related to this interpretation, Calvache and Willians (1992) and Banks *et al.* (1997) found that many of the pyroclastic flows deposits already Galeras volcano during the past 50 ka are lithic rich, with fines and clasts so strongly altered by hydrothermal action before eruption which evidences a long-lived hydrothermal system that indicates susceptibility to mass failure and explosive events.

Under the hypothesis that the spatial distribution of attenuation found in this work represents the combined effect of fractures that have been filled by hydrothermal material, we identify an important volume of magma at depth (> 13.0 km) that feeds minor shallow magma chambers throughout the fracture system. This fracture system plays a relevant role for the fumaroles activity and explosive eruptions. Stix *et al.* (1993) based on the record of explosive eruptions since May 1989 to Jun 1992 suggest that the actual magma under Galeras volcano is degassing at depth in the magma chamber and in the conduits. This degassing has reduced the H₂O, S, and Cl contents of the magma and generates a sealing process that forces gas-saturated magma to release volatile components into confined spaces, thereby causing pressurization and explosive eruptions. Stix *et al.* (1997) proposed a model of vulcanian eruption for Galeras assuming that two scenarios for pressurization related with six eruptions occurred during 1992-1993 involving edifice weakening

and the development of a shallow hydrothermal system from continued degassing of the magma plug. As weakening of the edifice formed an extensive system of fractures, the shallow hydrothermal system deepened due to meteoric water input and magmatic degassing. With time, a greater number of fractures were filled and partly sealed by hydrothermal minerals at progressively deeper levels. As a result, the source of the overpressure also deepened with time.

Similar models have been proposed by Fisher et al. (1996), Heiligmann et al. (1997), Alfaro and Zapata (1997) and Gil Cruz and Chouet (1997), who based on data of fumaroles, eruptive activity, diffuse degassing and long period seismic events identified that about some hundred meters at NW of the main crater is located the principal hydrothermal system which is structurally controlled. In this sector Ordoñez and Rey (1997) reported that at the end of 1990, a deformation source was located at a depth of 2-3 km. By December 1991, the source had migrated to less than 500 m in depth. Calvache and Williams (1997b) suggested that the dome formed was product of a magma mixing event occurred early in the history of its emplacement, and the uppermost part of the dome was degassing freely at the surface for ten months and produced an isolate the magmatic system. The July 1992 eruption tapped not only the uppermost part of the dome but also a deeper H₂O-saturated magma.

Calvache (1990) based on petrologic evidences, suggested a model to explain the eruptive processes at Galeras volcano. The model proposes the presence of a small magma reservoir located at little depth, which is connected to a deeper big deposit. Initial assumption of the model suggests that dacite magma suffers a fractioned crystallization process in the shallow chamber; latter, recharge events of basic material from the deep levels, generates mixture processes. Calvache concluded that disposition of a superficial chamber is responsible for causing vulcanian type explosions like consequence of mixture of shallow magma with deep basic magma. In agreement with this model, we can suggest that the main attenuation anomalies inverted from Q_c values correspond to the two magma deposits.

Moncayo et al. (2004) have reported temporal changes in the Q_c values at Galeras associated to changes in the volcanic activity. Along these changes they found a fall in the attenuation between 1989 and 1991, which has been associated with a degassing process that begins after the 1989 eruption. The rise between October 1991 and July 1992 was related to important changes in the volcanic activity, such as the extrusion of the andesitic dome with accompanying shallow deformation, increase in the rates and amplitudes of long-period seismic events, and a decrease in the SO₂ flux. Moncayo et al. (2004) suggested that the conduit was partially blocked because the viscous magma in the dome acted like a seal that stopped the gas flux, this caused a increase in the pressure and attenuation that ended with the July 1992 eruption. After this explosive eruption the dome was destroyed, and the gas released, so the pressure drops and so does the attenuation. A little rise took place before the June 1993 eruption, and they proposed that this can be caused by

the formation of small plugs that are destroyed in subsequent eruptions. In this scenario, the extension of the shallow attenuation anomaly from depth to surface is related with the presence of plugs and fluids that are constantly fed from a deeper body and produce the actual hydrothermal activity with eruptive events. The most recent event at November, 2005, generated the evacuation of people (aprox. 9000 persons) of the at least four towns around the Galeras volcano (Mapachico, Jenoy, La Florida and Nariño). However, the big volume involved in this process suggests that more eruptive events may be expected at the Galeras volcano.

6. Conclusion

We can conclude that the inversion process applied to estimate the spatial coda wave attenuation at Galeras volcano allowed us to determine the presence of two levels of anomalies which should be related to bodies of partially melted rocks that are located at 9.0 km and > 13.2 km from the summit. These bodies are responsible for causing vulcanian type explosions as a consequence of mixture of shallow magma with deep basic magma. The shallow body extends to the summit and represents the presence of plugs and fluids that are constantly fed from a deeper body and produce the actual hydrothermal activity with eruptive events.

Acknowledgements

We thank INGEOMINAS for the technical support for this study. We also thank colleagues from the Volcanological and Seismological Observatory of Pasto OVSP for their comments. This work has been partially founded by Geophysical Group and the Vicerrectoría Académica of the Universidad Nacional de Colombia.

References

- Aki, K. (1969). Analysis of the seismic coda of local earthquakes as scattered waves. *J. Geophys. Res.*, 74: 615-631.
- Aki, K., Chouet, B. (1975). Origin of coda waves: source, attenuation and scattering effects. *J. Geophys. Res.*, 80: 615-631.
- Alfaro, C. and Zapata J.A.† (1997). Acid gas emissions from Galeras Volcano, Colombia, 1989-1994. *Jour.Volcanol. Geotherm. Res.*, 77: 209-228.
- Banks, N., Calvache V., M. L. and Williams, S. N. (1997). Stratigraphy and radiometric ages of Volcán Galeras, Colombia. *Jour.Volcanol. Geotherm. Res.*, 77: 39- 56.
- Barrero, D. (1979). Geology of central western cordillera west of Buga and Roldanillo, Colombia. Internal report. INGEOMINAS, Bogotá (Colombia).
- Calvache, M. (1990). Geology and volcanology of the recent evolution of Galeras volcano,

- Colombia. M.S. Thesis, Louisiana State University.
- Calvache V., M. L. and Williams, S. N. (1992). Lithic-dominated pyroclastic flows at Galeras volcano, Colombia- An unrecognized volcanic hazard. *Geology*, 20 (6): 539-542.
- Calvache , M., Cortés, P., Williams, S. (1997). Stratigraphy and chronology of the Galeras Volcanic Complex, Colombia. *Jour.Volcanol. Geotherm. Res.*, 77: 5-19.
- Calvache, M.L. and Williams, S.N. (1997a). Geochemistry and petrology of the Galeras Volcanic Complex, Colombia. *Jour.Volcanol. Geotherm. Res.*,77: 21-38.
- Calvache, M.L. and Williams, S.N. (1997b). Emplacement and petrological evolution of the andesitic dome of Galeras volcano, 1990-1992. *Jour.Volcanol. Geotherm. Res.*,77: 57-69.
- Cepeda, H. (1986). Investigaciones petrológicas en el ambito de las planchas 429-Pasto y 410-La Union, con especial énfasis en el Complejo Volcánico del Galeras. Internal report. INGEOMINAS, Medellín (Colombia).
- Crosson, R.S. (1976). Crustal Structure Modeling of Earthquake Data, 1. Simultaneous Least Squares Estimation of Hypocenter and Velocity Partameters. *J. Geophys. Res.*, 81: 3036-3046.
- Fischer, T. P., Arehart, G. B., Sturchio, N. C., and Williams, S. N. (1996). The relationship between fumarole gas composition and eruptive activity at Galeras volcano, Colombia. *Geology*, 24 (6), 531- 534.
- Fischer, T. P., Arehart, G. B., Sturchio, N. C., Counce, D., Stix, J., and Williams, S. N. (1997). Chemical and Isotopic Composition of Fumarole Gases and Spring Discharges from Galeras Volcano, Colombia. *Jour.Volcanol. Geotherm. Res.*, 77: 229- 254.
- Gil Cruz, F. and Chouet B. (1997). Long-period events, the most characteristic seismicity accompanying the emplacement and extrusion of a lava dome in Galeras Volcano, Colombia, in 1991. *Jour.Volcanol. Geotherm. Res.*,77: 121-158.
- Gómez, D. M., and Torres, C., R.A. (1997). Unusual low-frequency volcanic seismic events with slowly decaying coda waves observed at Galeras and other volcanoes. *Jour.Volcanol. Geotherm. Res.*, 77: 173 -193
- Gómez, D. M., Torres, C., R.A., Seidl, D., Hellweg, M., and Rademacher H. (1999). Tornillo seismic events at Galeras volcano, Colombia: a summary and new information from broadband three-component measurements. *Annali di Geofisica*, 42(3): 365 - 378.
- Hurtado, A.D. and Cortés, G.P. (1997). Third version of the hazard map of Galeras Volcano, Colombia. *Jour.Volcanol. Geotherm. Res.*,77: 89-100.
- Heiligmann, M., Stix, J., Williams-Jones, G., Sherwood, B. and Garzón, G. (1997). Distal degassing of radon and carbon dioxide on Galeras volcano, Colombia. *Jour.Volcanol. Geotherm. Res.*,77: 267-283.
- Herrmann, R.B. (1980). Q estimates using the coda of local earthquakes. *Bull. Seism. Soc. Am.*, 70: 447-468.
- Hoshiba, M., 1991. Simulation of multiple scattered coda wave excitation based on the energy conservation law. *Phys. Earth Planet. Int.*, 67, 123-136.
- Lee, W. H., Lahr, J. C. (1975). HYPO71: A computer program for determining hypocenter,

- magnitude, and first motion pattern of local earthquakes. U.S. Geological Survey. Open-File report 75-311.
- Londoño, J. (1996). Temporal change in coda Q at Nevado del Ruiz Volcano, Colombia. *Jour.Volcanol. Geotherm. Res.*, 73: 129-139.
- Malin, P.E. (1978). A first order scattering solution for modeling lunar and terrestrial seismic coda, Ph.D. dissertation, Princeton University, Princeton New Jersey.
- Meissner, P.O., Flueh, E.R., Stribane, F. and Berg, E. (1976). Dynamics of the active boundary in southwest Colombia according to recent geophysical measurements. *Tectonophysics*, 35: 113-136.
- Moncayo, E. N., Vargas-Jiménez, C. A. and Duran-Tovar, J. P. (2004). Temporal variation of coda Q at Galeras volcano, Colombia. *Earth Sci. Res. J.*, 8 (1):19-24.
- Muñoz, F. A., Calvache, M. L., Cortés, G. P., Gomez, D. M., Narvaez, L., Ordonez, M., Ortega, A., Torres, R., Silva, B., Williams, S. N., Sanders, C. O., and Stix, J. (1993). Galeras Volcano, Colombia: the Decade Volcano Workshop and disastrous 14 January, 1993 eruption. *EOS*, 74 (26): 281- 287.
- Narváz, L., Torres, R., Gómez, D., Cortés, P., Cepeda, H., Stix, J. (1997). “Tornillo”-type seismic signals at Galeras volcano, Colombia, 1992–1993. *Jour.Volcanol. Geotherm. Res.*, 77: 159-171.
- Ordoñez, M. and Rey, C.A. (1997). Deformation associated with the extrusion of a dome at Galeras volcano, Colombia, 1990-1991. *Jour.Volcanol. Geotherm. Res.*,77: 115-120.
- Ortega, A., Torres, R., Gómez, D. (2002). Caracterización de fuentes sismogénicas de eventos volcano-tectónicos (VT) en el complejo volcánico Galeras. Paper presente at Primer Simposio Colombiano de Sismología, Bogotá, Colombia.
- Pulli, J. J. (1984). Attenuation of Coda Waves in New England. *Bull. Seism. Soc. Am.*, 74: 1149-1166.
- Rautian, T. J. and Khalturin, V.I. (1978). The Use of the Coda for the Determination of the Earthquake Source Spectrum. *Bull. Seismol. Soc. Am.* 68: 923 - 948.
- Sánchez, J.J., Gómez, D., Torres, R., Calvache, M.L., Ortega, A., Ponce, A.P., Acevedo, A.P., Gil Cruz, F., Londoño, J.M., Rodríguez, S.P., Patiño, J. and Bohorquez, P. (2005). Spatial mapping of the b-value at Galeras volcano, Colombia, using earthquakes recorded from 1995 to 2002. *Earth Sci. Res. J.*, 9(1): 30-36.
- Sano, Y., Gamo, T. and Williams, S. N. (1997). Secular Variations of Helium and Carbon Isotopes at Galeras Volcano, Colombia. *Jour.Volcanol. Geotherm. Res.*, 7: 255 - 266.
- Seidl, D., Hellweg, M., Rademacher, H., Gómez, M., D.M., and Torres, C., R.A. (1999). The anatomy of a tornillo: puzzles from three-component measurements at Galeras volcano (Colombia). *Annali di Geofisica*, 42 (3): 355 - 364.
- Stix, J., Zapata, J., Calvache, M., Cortez, G., Fischer, T., Gomez, D., Narvaez, L., Ordoñez, M., Ortega, A, Torres, C., Williams, S., (1993). A model of degassing at Galeras Volcano, Colombia, 1988-1993. *Geology*, 21: 963 - 967.
- Stix, J., Calvache V., M. L., and Williams, S. N., (1997). Galeras Volcanic Complex, Colombia, Introduction to an Interdisciplinary Study of a Decade Volcano. *Jour.Volcanol. Geotherm. Res.*, 77: 1- 4.

- Vargas, C. A., Ugalde, A., Pujades, L. And Canas, J. (2004). Spatial variation of coda wave attenuation in northwestern Colombia. *Geophys. J. Int.*, 158: 609-624.
- Ugalde, A., Vargas, C. A., Pujades, L. G. and Canas, J. A. (2002). Seismic coda attenuation after the Mw=6.2 Armenia (Colombia) earthquake of 25 January, 1999. *J. Geophys. Res.*, 107(B6), doi: 10.1029/2001JBO00197.
- Zapata, J., Calvache, M., Cortés, P., Fischer, T., Garzon, G., Gómez, D., Narváez L., Ordóñez, M., Ortega, A., Stix, J., Torres, R., Williams, S. (1997). SO₂ fluxes from Galeras Volcano, Colombia, 1989–1995: Progressive degassing and conduit obstruction of a Decade Volcano. *Jour.Volcanol. Geotherm. Res.*, 77: 195-208.